

# From Diapers to Decisions: MotherToBaby & NDBN Tackle Parents' Top Concerns

---

*By Lorrie Harris-Sagaribay, MPH, President of OTIS/MotherToBaby, in partnership with Joanne Samuel Goldblum, CEO and founder of National Diaper Bank Network.*

MotherToBaby is proud to partner with the **National Diaper Bank Network (NDBN)** as the organization marks its 15th anniversary in 2026 - celebrating a decade and a half of supporting families across the country by ensuring access to diapers and other essential material needs. Through our partnership, we've learned that many parents and caregivers, especially those who are pregnant or caring for newborns, often ask the same kinds of questions about everyday exposures during pregnancy and lactation.

These concerns come straight from the field, as NDBN staff and volunteers working directly with families hear questions like these every day. To help answer them, MotherToBaby experts weigh in on some of the families' most common questions about pregnancy and breastfeeding exposures.

First, it's important to remember that birth defects can happen in any pregnancy for different reasons. Out of all babies born each year, about 3 out of 100 (3%) will have a birth defect. Pregnancy problems (like miscarriage) can also happen in any pregnancy. MotherToBaby looks at research studies to understand if a certain exposure could increase the chance of birth defects or other problems in a pregnancy.

***Here are five of the exposures that NDBN families ask about most frequently:***

## 1. Caffeine

During pregnancy, it's recommended to keep your **caffeine** intake to less than 200 milligrams (mg) per day from all sources, including coffee, tea, energy drinks, and chocolate. (For reference, an 8-ounce cup of brewed coffee can contain 70 to 140 mg of caffeine, depending on the type of coffee and how it is brewed.) During breastfeeding, it's been suggested to stay under 300 mg per day, although that amount might be too high when the baby is very young (it takes young babies much longer than adults to process, or metabolize, caffeine). Even some older babies can be more sensitive to caffeine than others are. If you notice that your baby seems jittery, irritable, or wide awake in the middle of the night after you drink caffeine, you could consider cutting back.

Let's talk more about **energy drinks**. Most energy drinks contain not only caffeine, sugar, and B vitamins, but also **herbal ingredients** that most likely haven't been studied in pregnancy or breastfeeding. This means there is no information about how much of these ingredients might reach the developing baby during pregnancy or get into the breast milk. For this reason, it might be preferred to hold off on energy drinks until you aren't pregnant or breastfeeding any longer.

## 2. Over-the-Counter Pain Medications

Two of the most common over-the-counter pain medications are **acetaminophen** (such as Tylenol®) and **ibuprofen** (such as Advil®). These pain relievers are also found in multi-symptom products, such as cold medications. During pregnancy, using acetaminophen for a short time when directed by a healthcare provider to treat pain or fever has not been shown to increase pregnancy risks. Acetaminophen should be used as directed and only for as long as you need it to treat your condition. Ibuprofen, on the other hand, is typically not recommended in pregnancy, especially in the second half of pregnancy, unless your healthcare provider has specifically advised using it.

During breastfeeding though, acetaminophen and ibuprofen can both be used as directed without expected side effects for the baby. The amount of acetaminophen or ibuprofen that gets into the breast milk is much less than the dose that could be given directly to an infant.

## 3. Fish and Mercury

You might have heard a rumor that eating fish is not a good idea during pregnancy. But this is not the case. While it's true that most fish could contain some amount of mercury (or more specifically, a form of mercury called methylmercury), these amounts are often too small to increase pregnancy risks as long as you make thoughtful choices about eating fish.

The amount of **methylmercury in fish** gets higher as fish move up the food chain. Big predatory fish that eat smaller fish usually have higher levels of methylmercury. These big fish include swordfish, marlin, bigeye tuna, and king mackerel, among others. These fish should be avoided during pregnancy and while breastfeeding.

Other kinds of fish can be enjoyed in moderation (1 to 3 servings per week, depending on the kind of fish and where it is caught). The U.S. Food & Drug Administration (FDA) and United States Environmental Protection Agency (EPA) developed a helpful guide that lists many kinds of fish and gives advice on how often they can be eaten by women who are pregnant or breastfeeding and by children ages 1-11 years:  
<https://www.fda.gov/food/consumers/advice-about-eating-fish>.

## 4. Cleaning Products

Is it better to use natural cleaning products like vinegar during pregnancy or while breastfeeding? What about when you have a young child in the home?

The way you use a **cleaning product** is usually more important than what the product is. Although using vinegar as a cleaner isn't expected to increase risks during pregnancy or while breastfeeding, there could be some surfaces or situations that require more effective products for sanitizing and disinfecting. In order for a chemical to be able to reach a developing baby during pregnancy or to get into the breast milk, the chemical first has to be circulating in your bloodstream. As long as a product is used in a well-ventilated area (open doors or windows, turn on fans) and your skin is protected from direct contact with the cleaner, then using the cleaner as directed is unlikely to result in an exposure that would get into your bloodstream. If you start to feel any symptoms, such as nausea, dizziness, or headache, increase ventilation in the area and get some fresh air. If you do use vinegar to clean, be sure not to mix it with

ammonia or other chemicals, as that can create harmful fumes. With any cleaning product, follow the directions on the label for how to use and store it.

As far as what kinds of cleaners are preferred when you have young children in the home, that is a great question to talk about with your child's healthcare provider.

## 5. Animal Dander/Feces

Common household pets like dogs, cats, and rodents tend to be furry. Although pet dander itself is not known to be harmful during pregnancy or while breastfeeding, dander could increase the chance of breathing problems if you have significant allergies or **asthma**. If you are experiencing worsening symptoms of allergies or asthma during your pregnancy, be sure to talk about it with your healthcare provider. It's also a good idea to wash your hands after handling any kind of rodent, even sweet, furry, household pets.

What about animal feces? It's best to avoid direct contact with any animal feces during pregnancy. This means you get a free pass on cleaning the gerbil cage or scooping the litterbox, if possible. Cat feces, in particular, can contain a parasite responsible for an infection called **toxoplasmosis**, especially if the cat is allowed outside or is fed raw meat. A toxoplasmosis infection during pregnancy increases the chance of pregnancy complications, and could even pass to the fetus. There's no need to rehome your cat or avoid petting it, but you should avoid direct contact with the cat's feces while you are pregnant.

## Closing Thoughts

Through our partnership with NDBN, MotherToBaby is committed to answering the real questions families are asking—accurately, clearly, and compassionately. We know that parents want the best for their babies, and they deserve trusted, evidence-based information to make informed choices.

If you or someone you know has questions about exposures while pregnant or breastfeeding, you can contact a MotherToBaby specialist for free and confidential information via phone, text, or chat at **MotherToBaby.org**.

Together with NDBN, we'll continue listening, learning, and supporting families when it matters most.

## More About the National Diaper Bank Network

The National Diaper Bank Network (NDBN) leads a nationwide movement dedicated to helping individuals, children and families access the basic necessities they require to thrive and reach their full potential...including clean, dry diapers, period supplies and other basic needs. Launched in 2011 with the support of founding sponsor Huggies®, NDBN is dedicated to creating awareness of diaper need/diaper insecurity and advocating for public policy to end it. The Network is made up of more than 300 basic needs banks serving local communities throughout the U.S. More information on NDBN and diaper need is available at [nationaldiaperbanknetwork.org](https://nationaldiaperbanknetwork.org), and on **Twitter** (@DiaperNetwork), **Instagram** (@DiaperNetwork) and **Facebook** ([facebook.com/NationalDiaperBankNetwork](https://facebook.com/NationalDiaperBankNetwork)).

**Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://www.MotherToBaby.org).**

---

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, December 18, 2025.