

# Individualized Mental Health Care is Essential in Pregnancy

---

**Mental health conditions are real health issues that deserve appropriate and individualized medical care.** Making informed healthcare decisions requires that women and their healthcare providers be aware of the available safety information for each specific medication they might be considering. We encourage pregnant women and those planning a pregnancy to have open conversations with their healthcare providers to carefully weigh any potential increased risks of medications, as well as the risks of untreated or undertreated mental health conditions.

SSRIs – including commonly prescribed medications such as Zoloft (sertraline), Lexapro (escitalopram), Prozac (fluoxetine), Paxil (paroxetine), and others – are among the most well-studied medications used in pregnancy. Decades of research have shown that SSRIs, when used appropriately and under the care of a healthcare provider, can be part of an effective treatment plan for managing mental health conditions before, during, and after pregnancy, and can contribute to better outcomes for women and their babies.

As always, MotherToBaby and the Organization of Teratology Information Specialists (OTIS) stand with women and their healthcare teams by providing up-to-date, evidence-based information on the use of medications and other exposures during pregnancy and while breastfeeding.

MotherToBaby offers free, expert-reviewed fact sheets on SSRIs and other medications, as well as on conditions such as depression, anxiety, and stress. You can find many of our mental health resources here: <https://mothertobaby.org/pregnancy-breastfeeding-exposures/mental-health/>. Individuals can also reach out to a MotherToBaby specialist via phone, live chat, text, or email to receive personalized, confidential support.

---

**Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://mothertobaby.org).**

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, July 28, 2025.

---

# Individualized Mental Health Care is Essential in Pregnancy

---

Sara contacted us on the MotherToBaby live chat service after being referred by her OB/GYN who had told her that the Paxil (drug name paroxetine) she was taking could cause a heart defect. She was 14 weeks along and wanted to learn more, but was confused because all her genetic testing results had come back normal. She asked: “How could my baby have a heart defect if my non-invasive prenatal testing (NIPT) was negative?”

During pregnancy, there are many tests done to check on the health and development of your baby. It can be difficult to keep track of what they are all looking for and why they are important. To understand these tests, it can be helpful to know the differences between a birth defect and a chromosomal abnormality and what they each mean for the health of your baby.

All pregnancies have a background risk for birth defects of 3-5%, but certain exposures can increase the risk of birth defects above this background estimate. Most birth defects occur during the first trimester **while the baby is growing and developing**. A certain body part – such as the heart, brain, or limbs – might not develop correctly, and the baby’s body may look or function differently than it should. You may have heard of a baby being born with a hole in the heart, or with something called a cleft lip. These are two examples of birth defects that can occur during pregnancy. Birth defects can range from mild to severe, and the health of the baby will be dependent on where the birth defect occurs and how severe the problem is.

Certain tests done during pregnancy can look at your baby to see if there are birth defects. For example, most pregnant individuals will go in for an anatomy scan between 18 and 22 weeks where the healthcare provider will look at the baby using an ultrasound. Most people know this as the time when they can learn the gender of the baby, but the ultrasound will also take a detailed look at the baby’s organs and body parts, including the heart, brain, face, and stomach, to check for birth defects. While this is an important screening test, it is not perfect, and more minor defects may go unnoticed until birth.

Chromosomal abnormalities are changes in the baby’s DNA that happen at the time of conception. The best way to understand DNA is to think of it like a recipe book that holds all the recipes for the growth and development of different body parts. If a certain recipe calls for one stick of butter, but the printed book accidentally says two sticks of butter, the recipe will turn out different. In the same way, sometimes people have extra or missing amounts of DNA that can cause changes in development. For example, people who have a chromosomal abnormality known as Down syndrome have three copies of their 21st chromosome rather than only two. This extra amount of DNA is what causes the developmental differences in individuals with Down syndrome.

During your pregnancy, your healthcare provider may suggest that you meet with a genetic counselor to have non-invasive prenatal testing, or NIPT, performed. This test can be done as early as 10 weeks. During pregnancy, some of the baby’s DNA enters the pregnant woman’s bloodstream. By taking a small blood sample from mom, a lab can take a look at the baby’s DNA and tell if they have certain chromosomal abnormalities, such as Down syndrome. When you meet with a genetic counselor, they will go over all of the pros and cons of this test as well as your family history, and will meet with you again to review the results.

Back on the live chat service, Sara asked: “So because my NIPT results were normal, the baby is unlikely to have a chromosomal abnormality. However, a birth defect still could have happened in the first trimester, and I need to wait until my anatomy scan to get those results, is that right?” That’s exactly right, I replied. I then went on to explain that although Paxil (paroxetine) has been shown to increase the risk for heart defects in some studies, other studies do not suggest a risk, so the overall chance of the baby being affected is low.

As we ended our chat, Sara shared that she was feeling much more knowledgeable about the difference between a birth defect and a chromosomal abnormality. She was able to breathe a sigh of relief that her NIPT results came back normal, indicating a low risk for conditions like Down syndrome. She also felt much more confident heading to her anatomy scan in a few weeks knowing that this test, while not perfect, would be the best way to identify birth defects before her baby is born.

If you have any questions about birth defects or exposures during pregnancy, speak with a MotherToBaby specialist via phone, text, live chat, or email. For any questions regarding genetic testing in pregnancy, or to find a genetic counselor in your area, visit the Find a Genetic Counselor page on the National Society of Genetic Counselors website: <https://findageneticcounselor.nsgc.org/>

**Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://mother-to-baby.org).**

---

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, July 28, 2025.