

For Women With MS: Making Decisions About Pregnancy, Breastfeeding & More

By Elizabeth Salas, MPH, Teratology Information Specialist, MotherToBaby California

If you have Multiple Sclerosis (MS) and are currently pregnant, breastfeeding, or planning a pregnancy, where do you go when you have questions about MS or MS treatments? In this day and age, the first place you might go is the Internet. With no shortage of information at our fingertips, it may seem the answers to all of our questions are just a web search away. But when it comes to chronic conditions and treatments in pregnancy, reliable and accurate information isn't always easy to find, and the answers may not be so simple. So let's try a different approach, shall we? First, let's start with the facts!

The Good News

Studies about MS and pregnancy are encouraging. To date they show MS does not affect a woman's ability to get pregnant. For most women with MS, they are less likely to have a relapse during pregnancy especially in the 3rd trimester. Research shows pregnancy does not worsen MS or the progression of the disease¹. MS during pregnancy also does not increase the risk for birth defects, and does not increase the risk of major complications in pregnancy, during delivery, or for the newborn². In fact, some studies suggest pregnancy may have a protective effect for women, by slowing down or reducing the progression of MS – although more research is needed to confirm this finding³.

Making Progress in Treating MS

Ten years ago only a handful of treatments were available to treat MS. These medications, such as Betaseron® (Interferon Beta-1b) or Avonex® (Interferon Beta-1a), are called disease modifying medications because they slow down the natural course of MS while reducing the number and severity of relapses. Today there are twice as many disease modifying medications available – but the big question here is what do we know about these treatments during pregnancy or lactation?

The somewhat frustrating answer is that there is very little information about the safety of the newest medications during pregnancy or lactation. For this reason, standard practice has generally suggested women with MS stop treatment with disease modifying medications at least 1 menstrual cycle prior to attempting to conceive⁴. Older medications used to improve symptoms during a relapse, such as Solu-Medrol® (methylprednisolone) or prednisone, have been around since the 1950s⁵ and much has been published on their use in pregnancy⁶. (For more information about prednisone/prednisolone in pregnancy and lactation, see our **Fact Sheet**.) Regardless of which medications you may be taking, it's important for women with MS to plan their pregnancies and discuss treatment and options with your doctor before trying to become pregnant.

But what if your pregnancy, like nearly half of all pregnancies in the United States, was not planned? You may still have many questions about how your MS – and any medication you may be taking to treat it – could impact your pregnancy, such as: Could my medication have an effect on my developing baby? Will additional tests be needed during pregnancy to make sure my baby is all right? Should I continue taking my MS medications during pregnancy? If I stopped my medication, when can – or when should – I start taking them again? Can I breastfeed while taking these medications? The questions may seem overwhelming, but the good news is there are specialists who can answer your questions and they are just a phone call away!

Making The Call

Hello and thank you for calling MotherToBaby. We're here for you!

I am a counselor with MotherToBaby, a group of experts dedicated to providing women, healthcare providers, and the general public with accurate and up-to-date information on exposures during pregnancy and breastfeeding. We answer questions about everything from medications and cosmetics, to chronic conditions, like multiple sclerosis. Our service is FREE, confidential, and available for you. To speak to a counselor, call us toll free at (866) 626-6847.

Making A Difference

Every pregnant woman wants a healthy pregnancy. After personally talking to pregnant women with chronic conditions for nearly a decade, one thing has become very clear: we need better answers about how medications affect pregnancy. MotherToBaby has a follow-up program for pregnant women with MS, regardless of whether they are currently taking medication. We are learning more every day thanks to pregnant women with MS who are sharing information about their experiences. If you'd like to know more about current programs on MS and pregnancy, please contact one of our MotherToBaby Pregnancy Studies experts toll free at (877) 311-8972. You can help us make a difference, and together we can find the answers.



Elizabeth Salas is the Lead Teratology Information Specialist for MotherToBaby California, a non-profit that provides information to healthcare providers and the general public about medications and more during pregnancy and breastfeeding. She is based at the University of California, San Diego, and is passionate about the work MotherToBaby is doing to promote healthy moms, healthy pregnancies and healthy babies.

MotherToBaby is a service of the international Organization of Teratology Information Specialists (OTIS), a suggested resource by many agencies including the Centers for Disease Control and Prevention (CDC). If you have questions about medications, vaccines, diseases, or other exposures, call MotherToBaby toll-FREE at 866-626-6847 or call the Pregnancy Studies team directly at 877-311-8972. You can also visit MotherToBaby.org to browse a library of fact sheets, as well as visit our [Autoimmune Diseases and Pregnancy](#) page.

*Interested in more information about MS and pregnancy? Check out MotherToBaby's March 2014 blog, "**MS: The Diagnosis that Doesn't Mean Missing Out on Motherhood!**"*

References:

1. Baird, S. M., & Dalton, J. (2013). Multiple sclerosis in pregnancy. *Journal of Perinatal & Neonatal Nursing*, 27 (3), 232-41. doi: 10.1097/JPN.0b013e31829d98c5.
2. Tsui, A., & Lee, M. A. (2011). Multiple sclerosis and pregnancy. *Current Opinion in Obstetrics and Gynecology*, 23(6):435-9. doi: 10.1097/GCO.0b013e32834cef8f.
3. Roulet, E., Verdier-Taillefer, M. H., Amarenco, P., Gharbi, G., Alperovitch, A., & Marteau, R. (1993). Pregnancy and multiple sclerosis: a longitudinal Study of 125 remittent patients. *Journal of Neurology, Neurosurgery, & Psychiatry*, 56(10):1062-5.
4. Houtchens, M.K., & Kolb, C. M. (2013). Multiple sclerosis and pregnancy: therapeutic considerations. *Journal of Neurology*, 260(5):1202-14. doi: 10.1007/s00415-012-6653-9.

5. Clinical Pharmacology [database online]. Tampa, FL: Elsevier/Gold Standard, Inc.; 2014. URL: <http://www.clinicalpharmacology.com>. Updated August 2013 (Methylprednisolone) and September 2013 (Prednisone).

6. Briggs, G.G., Freeman, R. K., & Yaffe, S. J. (2011). *Drugs in Pregnancy and Lactation* (9th ed.). Philadelphia, PA: Lippincott Williams & Wilkins.

Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at MotherToBaby.org.

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, December 1, 2014.

For Women With MS: Making Decisions About Pregnancy, Breastfeeding & More

By Beth Kiernan, MPH, Interviewer & Teratogen Information Specialist, MotherToBaby

"I just found out I'm pregnant, and while I am glad, I'm also pretty overwhelmed," lamented Marian, a never-pregnant, newly-married woman who had struggled with Crohn's disease since her teenage years but was finally in semi-remission. "I'm not sure I'm comfortable taking my medications now, even though my doctors say they are fine. I can't even sleep! I am worried I'm harming my baby. I wonder if there will be long-term effects. I don't even know if it's OK to breastfeed and still take my medications, but I've had a terrible time in the past when my insurance changed and I didn't have access to these drugs."

"Your concerns are the concerns of every pregnant woman who takes medication for a chronic health condition," I told her gently. "This is the hallmark of motherhood, when a mother puts her baby's concerns above her own. However, in your case, you'll need to carefully consider the risks vs. the benefits of taking or not taking your medication. The goal is to keep your Crohn's disease from flaring so as to ensure as healthy a pregnancy as possible. Let me share some information with you. Once you have that knowledge, you will feel empowered, and you'll be able to sleep soundly knowing that you are making the best choices for both yourself and your baby."

How did Marian find me? I am a counselor at MotherToBaby Pregnancy Studies, conducted by the Organization of Teratology Information Specialists. Here at MotherToBaby, we help pregnant women, their partners, and their doctors by providing up-to-date information about exposures in pregnancy. What do we mean by "exposures"? Exposures

could include prescription medications, over the counter medications, herbal supplements, caffeine, alcohol, fever, hair dye...you name it, and the question has probably been asked and answered to the best of our ability by someone in our global network of counseling services.

Half of all pregnant women in the U.S. take at least one prescription medication in pregnancy.¹ For many of these women, taking their medication isn't a choice; it's vital to their health and well-being. I'm talking about women who take medications to control seizures, to treat psychiatric illness, to treat pain, to treat asthma, or to delay the progression of diseases such as Multiple Sclerosis or Rheumatoid Arthritis. Depending on the severity of your Inflammatory Bowel Disease (IBD), you may fall into this category too. It may be that NOT treating your Crohn's or Colitis will worsen your symptoms, and could lead to a situation where both you and your baby would be threatened. Lengthy hospitalization, severe pain, blood loss, surgery – all may outweigh the risk of taking the very medication that could prevent these grim developments.

And while many women plan ahead to get pregnant, consult with their doctors, consider their medication use in advance, and start or stop medications in anticipation of conception, over 50% of pregnancies in the U.S. are unplanned.² However pregnancy happens for you, it's important to learn about your medications and what others who take them have experienced with regard to pregnancy. Some medications were approved long ago by the Food and Drug Administration and we have years of good data on their use in pregnancy. Others, particularly newer biologics used to treat autoimmune diseases like IBD, may only have been on the market for a few years at the time you inquire, so there may not be a lot of information available about the use of the drug in pregnancy – which means you will need to weigh the risks of taking or not taking the medication to both you and your baby. Your doctor can help you assess those risks, and so can information specialists like me at MotherToBaby.

It will also be important to ask whether or not there could be any impact of your IBD medications to a breastfeeding baby. Undoubtedly, breastfeeding is far more beneficial to a baby than using commercial formula, but inquiring about actual infant experience with a mother's medication use will be valuable information for you and your baby. The American Academy of Pediatrics, and your pediatrician, can tell you if there are concerns for your newborn.

If you've given yourself the luxury of time by planning your pregnancy, speak with your gastroenterologist and your obstetrician or family doctor about your medications and your plans for a future pregnancy. Gather a team of experienced physicians who will support you in your choices, and avail yourself of their advice to ensure you will be healthy when the time comes to begin trying to get pregnant. Studies show that conceiving when your disease is in remission decreases your risk of having pregnancy complications and ongoing IBD flares.³ Also, now is a good time to begin taking a multivitamin or prenatal vitamin with folic acid to ensure the best start for your baby.

What if you aren't planning pregnancy this year? Then use this article as a starting point for considering the future. If you are of reproductive age, every sexual encounter presents the chance of becoming pregnant. It's important to know what you might face in terms of medication risks were you to unexpectedly become pregnant. Ask your prescribing doctor or the pharmacist about the research on your medication in pregnancy, or call MotherToBaby at our toll-free number 877-311-8972 and ask to speak with a counselor.



Beth Kiernan is a Teratogen Information Specialist with MotherToBaby Pregnancy Studies, a non-profit that conducts observational research about exposures in pregnancy and provides information to healthcare providers and the general public on medications and more during pregnancy and

breastfeeding. She is based at the University of California, San Diego, and is the married mother of four children ages 12 to 23 years.

MotherToBaby is a service of the international Organization of Teratology Information Specialists (OTIS), a suggested resource by many agencies including the Centers for Disease Control and Prevention (CDC). If you have questions about medications, vaccines, diseases, or other exposures, call MotherToBaby toll-FREE at 866-626-6847 or call the Pregnancy Studies team directly at 877-311-8972. You can also visit MotherToBaby.org to browse a library of fact sheets, including one on **IBD in Pregnancy, and find your nearest affiliate.**

References:

1. Mitchell, AA et. al. (2011). Medication use during pregnancy, with particular focus on prescription drugs: 1976-2008. *Am J Ob GYN*; 205(1): 51.e1-51.e8.
2. Finer LB and Zolna MR. (2011). Unintended pregnancy in the United States: incidence and disparities, 2006. *Contraception*; 84(5):478-485.
3. Abhyankar et. al. (2013). Meta-analysis: the impact of disease activity at conception on disease activity during pregnancy in patients with inflammatory bowel disease. *Aliment Pharmacol Ther.* 38(5):460-6.

Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at MotherToBaby.org.

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, December 1, 2014.

For Women With MS: Making Decisions About Pregnancy, Breastfeeding & More

By Neda Ebrahimi , Teratogen Information Specialist, Motherisk

As a counselor with **Motherisk**, the Canadian partner of MotherToBaby and a service of the Organization of Teratology Information Specialists (OTIS), I hear many stories from women about pregnancy. Some of those stories strike cords with me. Their urgency and desire to make the healthiest decisions possible for their future children is both

understandable and admirable. In honor of National Multiple Sclerosis Awareness Month, I give you **Nina's story**.

Nina's Story

"I'm 31 years old, and I was diagnosed with Relapsing Remitting Multiple Sclerosis (RRMS), when I was only 22. My first relapse was scary. I was writing my finals, and 2 days before my last final, I lost sight completely in one eye, and my legs felt so weak and wobbly that I couldn't stand even for a second. After going to the hospital and receiving several courses of steroids over 10 days, I started to improve but it took 2 months for my symptoms to fully resolve. And then, everything went back to normal, as if nothing had ever happened. I received my diagnosis several months after, and it felt like a death sentence. I had 2 more relapses before my doctor put me on disease modifying drug (DMD), and I started with Interferon-B1a. Over the last 8 years, I only experienced 5 more relapses. The last relapse I had was only a few months ago; I lost sight in my left eye, and numbness that ran from my face to my toes on just the right side of my body. I have always been able to work full-time except when I'm experiencing a relapse, for which I've had to take a month off. I am a dentist, so not surprisingly I can't carry out my job when I'm experiencing numbness in my hand. I met John 5 years ago at the MS clinic I used to visit. He was a nurse there. We fell in love, and despite of my illness he proposed to me last year, and we talked about having a family, with two children, hopefully one boy and one girl, and living happily ever after. It didn't initially worry me that one day I may want children. John is crazy about kids, and I feel my maternal instincts kick in every time I hold a baby. Since we got married, my anxiety has been increasing proportionally to my yearning for having a child. I know my MS can't be cured, at least not now, I know it can get worst over time, and eventually I may need support to carry out even simple tasks. Or Maybe I won't, and I would be one of the few who never enter the progressive state. I don't know if I'll be able to care for a baby and meet his or her demands. What will happen after my pregnancy? I really don't want to experience another relapse after I deliver. How am I going to manage my illness, and what will happen if I need to come off my DMD when I'm pregnant or breastfeeding? There are so many questions, and I don't know who to turn to."

Nina is not alone in her thirst for answers. MS is an autoimmune neurological disease with very different presentation. No two MS patients are exactly the same and symptoms can vary from just the occasional mild tingling in the finger tips to more severe symptoms that render the patient unable to walk or stand for several weeks. With Relapsing Remitting MS accounting for 85% of all MS cases, most patients will undergo a remissive state after an attack, and will resume their daily life with little or no hindrance. Some patients will continue to have modest symptoms during the remissive state which they learn to adapt to and manage by different medications and or lifestyle changes. As there are no current cures for MS, many MS patients live for decades with this disease, and must find the means to maintain a high quality of life as the disease progresses, which can be challenging in the later stages of the disease.

MS impacts many more women than men with a 3:1 ratio in North America. As the disease onset occurs during the reproductive ages, many women with MS face the dilemma of pregnancy at some point during their lives. Young women, like Nina, with MS planning pregnancies, have many questions. Because the disease presentation and progression varies from person to person, there is no exact answer and treatment and management must be tailored to the specific person's need. However, I'd like to address some of the most common questions to help all of the "Ninas" out there:

1. "Would the disease adversely impact the pregnancy and my developing baby?"

Up until the late 1950s, women with MS were advised to terminate their pregnancies. With our advancement in the field, we know that this is almost never necessary. Many women with MS continue to have healthy babies, and research shows that there is no increased risk for having a baby with a structural malformation or developmental delay and many deliver healthy babies with no major complications. Although there is a trend toward lighter weight babies, the birth weight percentile remains in the normal range for most. Another observation has been the higher rate of miscarriage in the MS population with mixed results from different studies. The reason for this is not well understood, but the majority of miscarriages are in early pregnancy. While miscarriage rates in the general population are around 10-15%, in women with MS the rates are closer to 20%-30%. With successful conception, the chance of delivering a healthy baby at term is high, and women with MS should be assured that their disease is unlikely to cause harm to the developing baby.

2. "Would my baby also have MS?"

There is a complex interplay between genetics and environment leading to MS. While the risk of getting MS in the general population is 0.3%, having a parent with MS will increase this risk by almost 15 times. So children of women

with MS may have a 3% to 6% chance of developing MS later in life, but the environmental and lifestyle factors may play the ultimate role in disease manifestation. Hence despite the genetic contribution, the risk for your baby developing MS remains small and can potentially be modified.

3. “If I stop my DMD when planning, what are the risks of having a relapse while I try to conceive?”

Depending on how long it takes to conceive, the drug free period prior to pregnancy may be a risky period for experiencing a relapse. While some women conceive after just one cycle, many will conceive after several months of actively trying to become pregnant. It will take 1 to 3 months (depending on the drug) to fully clear the system, and during this time, some may experience disease activity. If prior to starting the DMD you had very active disease, there is a risk that you’ll experience a relapse when you stop the medication, especially if it takes more than 3 months for you to conceive. The decision to continue DMDs is highly individualized and is determined on a case-by-case basis. You and your neurologist will determine the best mode of action.

4. Would having a pregnancy make my MS progress faster?

Pregnancy has not been shown to speed the disease process. In fact, pregnancy is a state of remission for many women with MS, and a time for optimal wellbeing. It is well established that relapse rates reduce by 70% by the third trimester of pregnancy compared to the year prior to pregnancy. However after delivery the relapse rate increases, with 60% of women experiencing a relapse in the first 3 to 6 months postpartum. While the risk is increased in the postpartum period, the course of MS tends to return to its baseline, and no worse than what it was in the year prior to pregnancy. Some studies have found a protective effect with pregnancy, with a delay in the long-term disease progression; however, more studies are needed to confirm this finding.

5. Would I be able to continue my DMD through the pregnancy?

Although many women with MS go through remission in the pregnancy, some will continue to experience disease activity especially in the first two trimesters. The decision to continue DMDs is dependent on several factors, including the type of medication, disease activity in the year prior to pregnancy, and the type of control achieved with the given DMD. The use of glatiramer, Interferon Beta 1a/1b, in pregnancy have not been associated with an increased risk for malformations and if you achieved great control with these drugs, and are at a high risk of relapsing, your physician may consider continuing your therapy through the pregnancy. The newer drugs, especially the oral DMDs, have not been well studied, therefore it is recommended that you discuss with your neurologist the best plan for the course of your pregnancy. There are ongoing research studies looking at the outcome of pregnancies following exposure to these medications. MotherToBaby and its affiliates are engaged in such studies. For study information or for the most up-to-date information about newer medications used to treat MS during pregnancy, call from anywhere in North America toll-FREE 866-626-6847.

6. What if I have a relapse during pregnancy?

While relapses during pregnancy are uncommon, they may happen, and can be quite severe for some women. Steroids are usually used to treat those relapses, although some success has been shown with IVIg therapy as well. A woman that experiences a severe debilitating relapse during her pregnancy, may require the standard steroid therapy, while a woman that experiences a mild flare-up may choose, in collaboration with her physician, to abstain from treatment. Systemic steroid use in the first trimester has been associated with a very small risk for cleft lip and palate, and use in the second half of pregnancy may increase the risk for having a smaller baby and for delivering prematurely (before 37 weeks gestation). However, it is recommended that you speak with your health care provider before you stop or change any medication. The benefits of taking a steroid and treating your condition should be weighed against these small possible risks. For more information, check out this fact sheet online:

<https://mothertobaby.org/fact-sheets/prednisoneprednisolone-pregnancy/> or call anywhere in North America toll-FREE 866-626-6847.

7. Should I breastfeed or start my DMD right after delivery?

The postpartum period is a period with a high risk of experiencing relapses. Data on whether breastfeeding has protective effect has conflicting results. Some studies suggest a protective effect, possibly due to the delay of menses returning, while others show no impact. Information on safety of DMDs in the breastfeeding period are scarce, however given the large molecule size of glatiramer acetate, and Interferons, it is unlikely any will transfer into milk. If they do, they are likely not to be absorbed from the baby’s gastrointestinal tract. There is no information regarding other DMD usages during lactation. The benefits of breastfeeding baby are numerous, but, ultimately, your functionality and ability to care for your child take priority. The decision to breastfeed or not may depend on your ability to breastfeed, especially since the demands of a newborn and the hormonal changes in the postpartum period can be very taxing on your energy levels and if you experience chronic fatigue due to your condition. Thus, if a woman (while consulting her physician) decides to breastfeed she may do so. However, if she needs to restart her DMD, currently she may be

advised to stop breastfeeding.

Bottomline: While having MS poses physical and emotional challenges, it does not jeopardize a woman's capacity to motherhood. With careful planning and close collaboration with your doctors and healthcare providers, and especially with some support from family and friends, you will be able to have successful pregnancies, healthy children, and out of control teenagers, just like any other woman. So if becoming a mother is something you have always wanted and looked forward to, having MS is more of a bump in the road rather than a life sentence, and with some maneuvering you can achieve your dreams. Happy parenthood!

Neda Ebrahimi is a research associate and counselor at the Canadian Motherisk program, a non-profit MotherToBaby/OTIS affiliate that aims to educate the public about medications and more during pregnancy and breastfeeding. The Motherisk program is also a center for teaching and clinical research in the area of exposures in pregnancy and breastfeeding. Neda is pursuing her PhD in the field of Multiple Sclerosis in Pregnancy. To learn more about her work and about her study, email her at neda.ebrahimi@sickkids.ca or call 416-813-7654 ext. 204928. You can also call the Motherisk Helpline at 1-877-439-2744 and ask to be referred to the MS study.

MotherToBaby and its affiliates are services of the international Organization of Teratology Information Specialists (OTIS), a suggested resource by many agencies including the Centers for Disease Control and Prevention (CDC). If you have questions about MS, medications or other exposures during pregnancy or breastfeeding, call toll-FREE 866-626-6847 or visit MotherToBaby.org.

Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at MotherToBaby.org.

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, December 1, 2014.