

# Lidocaine

---

## Selected References:

- Abboud TK, et al. 1983. Lack of adverse neonatal neurobehavioral effects of lidocaine. *Anesth Analg*; 62:473-475.
- Abboud TK, et al. 1984. Continuous infusion epidural analgesia in parturients receiving bupivacaine, chloroprocaine, or lidocaine – maternal, fetal, and neonatal effects. *Anesth Analg*; 63:421-428.
- Actavis Pharma, Inc. 2019. Lidocaine and Prilocaine Cream Drug Label. Available at: <https://dailymed.nlm.nih.gov/dailymed/druginfo.cfm?setid=1972d657-2d5a-4697-bba9-80caffc2f2d7>. Accessed 29 July 2025.
- Baradari AG, et al. 2017. Bolus administration of intravenous lidocaine reduces pain after an elective caesarean section: Findings from a randomised, double-blind, placebo-controlled trial. *J Obstet Gynaecol*. 37(5):566-570.
- da Cunha YGM et al. 2025. The Use of Different Local Anesthetics in Pregnant Women in Dentistry: A Systematic Review. *Curr Rev Clin Exp Pharmacol*. Apr 21. doi: 10.2174/0127724328349965250407082245. Online ahead of print.
- Demeulemeester V, et al. 2018. Transplacental lidocaine intoxication. *J Neonatal-Perinatal Med*; 11:439-441.
- Dryden RM, Lo MW. 2000. Breast milk lidocaine levels in tumescent liposuction. *Plast Reconstr Surg*; 105:2267-2268.
- Favero V, et al. 2021. Pregnancy and Dentistry: A Literature Review on Risk Management during Dental Surgical Procedures. *Dent J (Basel)*. 9(4):46.
- Fujinaga M, Mazzeo RI. 1986. Reproductive and teratogenic effects of lidocaine in Sprague-Dawley rats. *Anesthesiology*. 65:626-632.
- Heinonen OP, et al. 1977. *Birth Defects and Drugs in Pregnancy*. Publishing Sci Group, Littleton, MA.
- Kuhnert BR, et al. 1984. Effects of maternal epidural anesthesia on neonatal behavior. *Anesth Analg*; 63:301-308.
- Kuhnert BR, et al. 1986. Lidocaine disposition in mother, fetus, and neonate after spinal anesthesia. *Anesth Analg* 65:139-144.
- Lebedevs TH, et al. 1993. Excretion of lignocaine and its metabolite monoethylglycinexylidide in breast milk following its use in a dental procedure. A case report. *J Clin Periodontol* 20: 606-608.
- Li JE, et al. 2019. Cutaneous Surgery in Patients Who Are Pregnant or Breastfeeding. *Dermatol Clin*. 37(3):307-317. Murzaku EC, et al. 2021. Surgical management and practices in pregnancy and lactation: A survey of United States dermatologic surgeons. *J Am Acad Dermatol*. 84(4):1134-1136.
- Ortega D, et al. 1999. Excretion of lidocaine and bupivacaine in breast milk following epidural anesthesia for cesarean delivery. *Acta Anaesthesiol Scand*; 43:394-397.
- Silveira MPT, et al. 2020. Breastfeeding and risk classification of medications used during hospitalization for delivery: 2015 Pelotas Birth Cohort. *Rev Bras Epidemiol*. 23:e200026.
- Watson PD, Ott MA. 1982. Lidocaine and mepivacaine in cord blood. *Ped Pharm*; 2:341-348.
- Wikland, M., Evers, H., Jakobsson, A.-H., Sandqvist, U., & Sjöblom, P. (1990). The concentration of lidocaine in follicular fluid when used for paracervical block in a human IVF-ET programme. *Human Reproduction*, 5(8), 920-923
- Zeisler JA, et al. 1986. Lidocaine excretion in breast milk. *Drug Intell Clin Pharm*; 20:691-693.

Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://www.MotherToBaby.org).

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, August 1, 2025.

---

# Lidocaine

---

## Selected References:

- Dentsply Pharmaceutical, Inc. 2021. Prilocaine injection drug label. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=db23a56f-1e41-4843-9220-1b2e3059db41>. [Accessed 8/2025].
- Fougera & Co. 2024. Lidocaine and prilocaine cream drug label. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=833cd52c-6470-49c4-937f-1393971f4db9>. [Accessed 8/2025].
- Erol S, et al. 2017. Transient methemoglobinemia in three neonates due to maternal pudendal anesthesia. J Coll Physicians Surg Pak. 27(12):783-784.
- Guay J. 2009. Methemoglobinemia related to local anesthetics: a summary of 242 episodes. Anesth Analg. 108(3):837-45.
- Kirschbaum M, et. al. 1991. [Fetal methemoglobinemia caused by prilocaine—is use of prilocaine for pudendal block still justified?]. Geburtshilfe Frauenheilkd. 51(3):228-30.
- Uslu S, Comert S. 2013. Transient neonatal methemoglobinemia caused by maternal pudendal anesthesia in delivery with prilocaine: report of two cases. Minerva Pediatr. 65(2):213-7.

Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://www.MotherToBaby.org).

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, August 1, 2025.

---

# Lidocaine

---

## Selected References:

- Gilboa SM, et al. 2009. National Birth Defects Prevention Study: Use of antihistamine medications during early pregnancy and isolated major malformations. *Birth Defects Res A Clin Mol Teratol* 85(2):137-150.
- Gilboa SM, et al. 2014. Antihistamines and birth defects: a systematic review of the literature. *Expert opinion on drug safety*. 13.12: 1667-98.
- Hansen Craig, et al. 2020. Use of antihistamine medications during early pregnancy and selected birth defects: The National Birth Defects Prevention Study, 1997-2011. *Birth defects research*16: 1234-52.
- Ito S, et al. 1993. Prospective follow-up of adverse reactions in breast-fed infants exposed to maternal medication. *Am J Obstet Gynecol* 168:1393-9.
- Kallen B, Mottet I. 2003. Delivery outcome after the use of meclozine in early pregnancy. *Eur J Epidemiol*. 18:665-669.
- Katselou M, et al. 2018. A fully validated method for the simultaneous determination of 11 antihistamines in breast milk by gas chromatography-mass spectrometry. *Biomed Chromatogr* 32(8):e4260.
- Lenz W. 1966. Malformations caused by drugs in pregnancy. *Am J Dis Child* 112:99-106.
- Messinis IE, et al. 1985. Histamine H1 receptor participation in the control of prolactin secretion in postpartum. *J Endocrinol Investig* 8:143-6.
- Michaelis J, et al. 1983. Prospective study of suspected associations between certain drugs administered during early pregnancy and congenital malformations. *Teratology* 27:57-64.
- Mondillo C, et al. 2018. Potential negative effects of anti-histamines on male reproductive function. *Reproduction*. 155.5: R221-7.

Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://www.MotherToBaby.org).

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, August 1, 2025.

---

# Lidocaine

---

## Selected References:

- Ceyhan ST, et al. 2010. Serum vitamin B12 and homocysteine levels in pregnant women with neural tube defect. *Gynecol Endocrinol.* 26(8):578-581.
- Dror DK, Allen LH. 2018. Vitamin B-12 in human milk: A systematic review. *Adv Nutr* 9:358s-66s.
- Duggan C, et al. 2014. Vitamin B-12 supplementation during pregnancy and early lactation increases maternal, breast milk, and infant measures of vitamin B-12 status. *J Nutr.* 144(5):758-764.
- Groenen PMW, et al. 2004. Marginal maternal vitamin B12 status increases the risk of offspring with spina bifida. *Am J Obstet Gynecol* 191(1):11-17.
- Hampel D, Allen LH. 2016. Analyzing B-vitamins in human milk: Methodological approaches. *Crit Rev Food Sci Nutr* 56:494-511.
- Lai JS, et al. 2019. Maternal plasma vitamin B12 concentrations during pregnancy and infant cognitive outcomes at 2 years of age. *Br J Nutr.* 121(11):1303-1312.
- Munger RG, et al. 2021. Maternal vitamin B12 status and risk of cleft lip and cleft palate birth defects in Tamil Nadu State, India. *Cleft Palate Craniofac J* 58(5):567-576.
- National Institute of Health Office of Dietary Supplements. 2024. Vitamin B12: Fact Sheet for Health Professionals. Available at: <https://ods.od.nih.gov/factsheets/VitaminB12-HealthProfessional/>. Accessed 6 Jun 2024.
- Nelen WL, et al. 2000. Hyperhomocysteinemia and recurrent early pregnancy loss: a meta-analysis. *Fertil Steril.* 74(6):1196-1199.
- Ray JG, Blom HJ. 2003. Vitamin B12 insufficiency and the risk of fetal neural tube defects. *QJM.* 96(4):289-295.
- Ray JG, et al. 2007. Vitamin B12 and the risk of neural tube defects in a folic-acid-fortified population. *Epidemiology* 18(3):362-366.
- Reznikoff-Etievant MF, et al. 2002. Low Vitamin B(12) level as a risk factor for very early recurrent abortion. *Eur J Obstet Gynecol Reprod Biol.* 104(2):156-159.
- Senousy SM, et al. 2018. Association between biomarkers of vitamin B12 status and the risk of neural tube defects. *J Obstet Gynaecol Res.* 44(10):1902-1908.
- Suarez L, et al. 2003. Maternal serum B12 levels and risk for neural tube defects in a Texas-Mexico border population. *Ann Epidemiol* 13(2):81-88.
- Van Rooij IALM, et al. 2003. Vitamin and homocysteine status of mothers and infants and the risk of nonsyndromic orofacial clefts. *Am J Obstet Gynecol* 189(4):1155-1160.
- Wilson A, et al. 1999. A common variant in methionine synthase reductase combined with low cobalamin (vitamin B12) increases risk for spina bifida. *Mol Genet Metab.* 67(4):317-323.
- Zhang T, et al. 2009. Maternal serum vitamin B12, folate and homocysteine and the risk of neural tube defects in the offspring in a high-risk area of China. *Public Health Nutr.* 12(5):680-686.

Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://www.MotherToBaby.org).

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, August 1, 2025.

---

# Lidocaine

---

## Selected References:

- Al-Saleh I, et al. 2020. Effects of early and recent mercury and lead exposure on the neurodevelopment of children with elevated mercury and/or developmental delays during lactation: A follow-up study. *Int J Hyg Environ Health* 230:113629.
- Ashrap P, et al. 2020. Maternal blood and metalloids concentrations in association with birth outcomes in Northern Puerto Rico. *Environ Int* 138:105606.
- Buck Louis GM, et al. 2017. Low-level environmental metals and metalloids and incident pregnancy loss. *Reprod Toxicol* 69:68-74.
- Byeong-Jin Y, et al. 2016. Evaluation of mercury exposure level, clinical diagnosis and treatment for mercury intoxication. *Ann Occup Environ Med* 28:5.
- Choy CMY, et al. 2002. Relationship between semen parameters and mercury concentration in blood and in seminal fluid from subfertile males in Hong Kong. *Fertil Steril* 78(2):426-428.
- Cox C, et al. 199. Prenatal and postnatal methyl mercury exposure and neurodevelopmental outcomes. *JAMA* 282:1333-1334.
- Crump KS, et al. 1998. Influence of prenatal mercury exposure upon scholastic and psychological test performance: Benchmark analysis of a New Zealand cohort. *Risk Anal* 18:701-713.
- Davidson PW, et al. 1998. Effects of prenatal and postnatal methyl mercury exposure from fish consumption on neurodevelopment. *JAMA* 280:701-707.
- Davidson PW, et al. 2006. Prenatal methylmercury exposure from fish consumption and child development: A review of evidence and perspectives from the Seychelles Child Development Study. *Neurotoxicology* 27:1106-1109.

- Debes F, et al. 2016. Cognitive deficits at age 22 years associated with prenatal exposure to methylmercury. *Cortex* 74:358-69.
- Dorea JG, 2004. Mercury and Lead during breast-feeding. *British J of Nutr* 92(1):21-40.
- Dorea JG. 2021. Exposure to environmental neurotoxic substances and neurodevelopment in children from Latin America and the Caribbean. *Environ Res* 192:110199.
- Food and Drug Administration (FDA). 2017. Mercury levels in commercial fish and shellfish (1990–2012). Available at: <https://www.fda.gov/food/foodborneillnesscontaminants/metals/ucm115644.htm>
- Golding J, et al. 2017. Maternal prenatal blood mercury is not adversely associated with offspring IQ at 8 years provided the mother eats fish: A British prebirth cohort study. *Int J Hyg Environ Health* 220(7): 1161-1167.
- Golding J, et al. 2022. The benefits of fish intake: Results concerning prenatal mercury exposure and child outcomes from the ALSPAC prebirth cohort. *Neurotoxicology* 91:22-30.
- Gokoel AR, et al. 2020. Influence of prenatal exposure to mercury, perceived stress and depression on birth outcomes in Suriname: Results from the MeKiTamara Study. *Int J Environ Res Public Health* 17(12):4444.
- Grandjean P, et al. 1997. Cognitive deficit in 7-year-old children with prenatal exposure to methyl mercury. *Neurotoxicol Teratol* 19:417-428.
- Grandjean P, et al. 1999. Methylmercury exposure biomarkers as indicators of neurotoxicity in children aged 7 years. *Am J Epidemiol* 149:301-305.
- Grandjean P, et al. 2003. Attenuated growth of breast-fed children exposed to increased concentrations of methylmercury and polychlorinated biphenyls. *FASEB J* 17(6):699-701.
- Hibbeln J, et al. 2018. Total mercury exposure in early pregnancy has no adverse association with scholastic ability of the offspring particularly if the mother eats fish. *Environ Int* 116:108-115.
- Hibbeln JR, et al. 2019. Relationship between seafood consumption during pregnancy and childhood neurocognitive development: Two systematic reviews. *Prostaglandins Leuko Essent Fatty Acids* 151:14-36.
- Howe CG, et al. 2021. Prenatal metal mixtures and birth weight for gestational age in a predominately lower-income Hispanic pregnancy cohort in Los Angeles. *Environ Health Perspect* 128(11):117001.
- Hsi HC, et al. 2014. The neurological effects of prenatal and postnatal mercury/methylmercury exposure on three-year-old children in Taiwan. *Chemosphere* 100:71-76.
- Hu Y, et al. 2016. Prenatal low-level mercury exposure and infant neurodevelopment at 12 months in rural northern China. *Environ Sci Pollut Res Int* 23(12): 12050-12059.
- Iwai-Shimada M, et al. 2015. Methylmercury in the breast milk of Japanese mothers and lactational exposure of their infants. *Chemosphere*. 126:67-72.
- Jensen TK, et al. 2005. Effects of breast feeding on neuropsychological development in a community with methylmercury exposure from seafood. *J Expo Anal Environ Epidemiol* Sep;15(5):423-430.
- Kim B, et al. 2020. Adverse effects of prenatal mercury exposure on neurodevelopment during first 3 years of life modified by early growth velocity and prenatal maternal folate level. *Environ Res* 191:109909.
- Klus JK, et al. 2023. Postnatal methylmercury exposure and neurodevelopmental outcomes at 7 years of age in the Seychelles Child Development Study Nutrition Cohort 2, *NeuroToxicology*, 99: 115-119,
- Koos BJ & Longo LD. 1976. Mercury toxicity in the pregnant woman, fetus, and newborn infant. *Am J Obstet Gynecol* 390(5):390-409.
- Lamoureux-Tremblay V, et al. 2021. Altered functional activations of prefrontal brain areas during emotional processing of fear in Inuit adolescents exposed to environmental contaminants. *Nurotoxicol Teratol* 85:106973.
- Myers GJ, et al. 2003. Prenatal methyl mercury exposure from ocean fish consumption in the Seychelles child development study. *Lancet* 361:1686-1692.
- Myers, et al. 2009. Postnatal exposure to methyl mercury from fish consumption: a review and new data from the Seychelles Child Development Study. *Neurotoxicology*, 30(3):338-349
- Myers GJ, et al. 2007. Nutrient and methyl mercury exposure from consuming fish. *J Nutr* 137(12):2805-2808.
- Minguez-Alarcon L, et al; Earth Study Team. 2017. Hair mercury (Hg) levels, fish consumption and semen parameters among men attending a fertility center. *Int J Hyg Environ Health* pii:S1438-4639(17)30315-2.
- Nišević RJ, et al. 2019. Combined prenatal exposure to mercury and LCUPUFA on newborn's brain measures and

neurodevelopment at the age of 18 months. *Environ Res* 178:108682.

- Oken E and Bellinger DC. 2008. Fish consumption, methylmercury and child neurodevelopment. *Curr Opin Pediatr* 20(2):178-183.
- Papadopoulou E, et al. 2021. Maternal seafood intake during pregnancy, prenatal mercury exposure and child body mass index trajectories up to 8 years. *Int J Epidemiol* 1-13. Online ahead of print.
- Polevoy C, et al. 2020. Prenatal exposure to legacy contaminants and visual acuity in Canadian infants: a maternal-infant research on environmental chemicals study. *Environ Health* 19(1):14.
- Rothenberg SE, et al. 2021. Maternal methylmercury exposure through rice ingestion and child neurodevelopment in the first three years; a prospective cohort study in rural China. *Environ Health* 20(1):50.
- Saavedra S, et al. 2021. Impact of dietary mercury intake during pregnancy on the health of neonates and children: a systematic review. *Nut Rev* nuab029
- Saito, H. 2020 (Reproduction from 2004). Congenital Minamata disease: a description of two cases in Niigata. *Neurotoxicology* 81:360-363.
- Schaefer C, et al 2007. Industrial Chemicals and environmental contaminants: Mercury ***In Drugs During Pregnancy and Lactation***, pgs 816-817. Amsterdam
- Shamlaye C, et al. 2020. The Seychelles Child Development Study: Two decades of collaboration. *Reproduction in Neurotoxicology*. 81:315-322.
- Skerfving & Copplestone. 1976. Poisoning caused by the consumption of organomercury-dressed seed in Iraq. *Bull World Health Organ* 1976;54(1):101-112.
- Sloane-Reeves J, et al. 2020. Scholastic achievement among children enrolled in the Seychelles Child Development Study. *Neurotoxicology* 81:347-352.
- Smith JC, et al. 1997. Hair methylmercury levels in U.S. women. *Arch Environ Health*. 52(6):476-80.
- Spiller P, et al. 2019. An abundance of seafood consumption studies presents new opportunities to evaluate effects on neurocognitive development. *Prostaglandins Leukot Essent Fatty Acids* 151:8-13.
- Strain J. 2014. Eating fish for two. *Nutr Bull* 39(2):181-186.
- Strain JJ, et al. 2015. Prenatal exposure to methyl mercury from fish consumption and polyunsaturated fatty acids: associations with child development at 20 mo of age in an observational study in the Republic of Seychelles. *Am J Clin Nutr* 101(3):530-537.
- Strain JJ, et al. 2021. Associations of prenatal methylmercury exposure and maternal polyunsaturated fatty acid status with neurodevelopmental outcomes at 7 years of age: results from the Seychelles Child Development Study Nutrition Cohort 2. *Am J Clin Nutr* 113(2):304-313.
- Tian T, et al. 2021. Single and mixed effects of metallic elements in maternal serum during pregnancy on risk for fetal neural tube defects: A Bayesian kernel regression approach. *Environ Pollut* 285:117203.
- Tong M, et al. 2021. Total mercury concentration in placental tissue, a good biomarker of prenatal mercury exposure, is associated with risk for neural tube defects in offspring. *Environ Int* 150:106425.
- Ulloa AC, et al. 2021. Prenatal methylmercury exposure and DNA methylation in seven-year-old children in the Seychelles Child Development Study. *Environ Int* 147:106321.
- US DHHS: Mercury Toxicity, Monograph 17. ATSDR Case Studies in Environmental Medicine, 1992.
- United States Environmental Protection Agency (EPA). Choose fish and shellfish wisely. <https://www.epa.gov/choose-fish-and-shellfish-wisely>.
- United States Environmental Protection Agency (EPA). 2025. Choose Fish and Shellfish Wisely. <https://www.epa.gov/choose-fish-and-shellfish-wisely>.
- United States Environmental Protection Agency (EPA). Estimated Fish Consumption Rates for the U.S. Population and Selected Subpopulations (NHANES 2003-2010). Final Report. April 2014. EPA-820-R-14-002. <https://www.epa.gov/sites/production/files/2015-01/documents/fish-consumption-rates-2014.pdf>.
- United States Environmental Protection Agency (EPA). Trends in Blood Mercury Concentrations and Fish Consumption Among U.S. Women of Reproductive Age (EPA NHANES, July 2013).
- United States Environmental Protection Agency (EPA) 2025. National Biomonitoring Program. Mercury Factsheet. <https://www.epa.gov/americaschildrenenvironment/biomonitoring-mercury> [Accessed 2025].

- United States Food and Drug Administration (FDA). Eating Fish for Those Who Might Become or Are Pregnant or Breastfeeding and Children Ages 1 to 11 Years. <https://www.fda.gov/food/consumers/questions-answers-fdaepa-advice-about-eating-fish-those-who-might-become-or-are-pregnant-or> [Accessed August 2025].
- United States Food and Drug Administration (FDA) & United States Environmental Protection Agency (EPA). Eating Fish: What Pregnant Women and Parents Should Know. <https://www.fda.gov/food/consumers/advice-about-eating-fish> [Accessed August 2025].
- van Wijngaarden E, et al. 2017. Methyl mercury exposure and neurodevelopmental outcomes in the Seychelles Child Development study main cohort at age 22 and 24 years. *Neurotoxicol Teratol* 59:35-42.
- Vigeh M, et al. 2018. Prenatal mercury exposure and birth weight. *Reprod Toxicol* 76:78-83.
- Weichselbaum E, et al. 2013. Fish in the diet: a review. *Nutrition Bulletin* 38: 128-177.
- Xiang H, et al. 2019. Protective effect of high zinc levels on preterm birth induced by mercury exposure during pregnancy: A birth cohort study in China. *J Trace Elem Med Biol* 55:71-77.
- Young EC, et al. 2020 (Reproduction from 2004). Association between prenatal dietary methyl mercury and developmental outcomes on acquisition of articulatory-phonologic skills in children in the Republic of Seychelles. *Neurotoxicology* 81:353-357.

**Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://www.MotherToBaby.org).**

---

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, August 1, 2025.