

Dextroamphetamine

This sheet is about exposure to dextroamphetamine in pregnancy and while breastfeeding. This information should not take the place of medical care and advice from your healthcare provider.

What is dextroamphetamine?

Dextroamphetamine or d-amphetamine is a prescription medication used for attention deficit hyperactivity disorder (ADHD) and narcolepsy (extreme daytime sleepiness). It has also been prescribed to treat some mental health conditions. Some brand names include Dexedrine®, Dexedrine Spansules®, Dextrostat®, Liquadd®, ProCentra®, and Zenzedi®. Dextroamphetamine is combined with amphetamine to make Adderall®. Dextroamphetamine is an amphetamine and can become addictive.

Sometimes when people find out they are pregnant, they think about changing how they take their medication, or stopping their medication altogether. However, it is important to talk with your healthcare providers before making any changes to how you take this medication. Stopping dextroamphetamine suddenly can cause withdrawal in some people. If you are going to stop using this medication, your healthcare providers might talk with you about slowly reducing your dose over time. Your healthcare providers can also talk with you about the benefits of treating your condition and the risks of untreated illness during pregnancy.

I take dextroamphetamine. Can it make it harder for me to become pregnant?

Dextroamphetamine has not been studied to see if using it could make it harder to get pregnant.

Does taking dextroamphetamine increase the chance of miscarriage?

Miscarriage can occur in any pregnancy. Studies have not been done to see if dextroamphetamine increases the chance for miscarriage.

Does taking dextroamphetamine increase the chance of birth defects?

Every pregnancy starts out with a 3-5% chance of having a birth defect. This is called the background risk. Using dextroamphetamine as prescribed for medical reasons does not appear to increase the chance of birth defects. Misuse of dextroamphetamine (taking more than prescribed amounts) has not been well studied.

Does taking dextroamphetamine in pregnancy increase the chance of other pregnancy related problems?

One older study found a small effect on birth weight among people who continued to use prescribed dextroamphetamine past 28 weeks of pregnancy to control weight gain, but other studies looking at the use of dextroamphetamine with amphetamine to treat ADHD have not found an effect on birth weight.

Research shows that misuse of amphetamines (using more than prescribed) may increase the chance for poor growth (babies born small and/or with a small head size), low birth weight (weighing less than 5 pounds, 8 ounces [2500 grams] at birth), and preterm delivery (birth before week 37). The pregnancies in these studies may have had other risk factors, such as exposure to alcohol and cigarettes, or not having prenatal care.

I need to take dextroamphetamine throughout my entire pregnancy. Will it cause withdrawal symptoms in my baby after birth?

The misuse of other amphetamines (such as methamphetamine) during pregnancy can cause temporary symptoms in newborns soon after birth. These symptoms are sometimes referred to as withdrawal. They can include jitteriness, sleepiness, and trouble breathing at the time of birth. These symptoms have not been reported in the newborns of people who are taking dextroamphetamine as prescribed by their healthcare providers.

Does taking dextroamphetamine in pregnancy affect future behavior or learning for the child?

Studies have not been done to see if dextroamphetamine can cause behavior or learning issues for the child.

Breastfeeding while taking dextroamphetamine:

When dextroamphetamine is taken as directed, it gets into breastmilk in low levels. Babies can be monitored for any symptoms related to dextroamphetamine, especially if they are born preterm (before 37 weeks) or are under two months of age. If you suspect the baby has any symptoms such as decreased appetite, sleeplessness, or irritability, contact the child's healthcare provider. A small study of four infants whose mothers were taking dextroamphetamine for ADHD found no problems in the health and growth of those infants up to 6-10 months of age. Some evidence suggests that large doses of dextroamphetamine could lower milk supply in people who are newly breastfeeding.

The product labels for some dextroamphetamine products recommend people who are breastfeeding not use these medications. But, the benefit of using dextroamphetamine may outweigh possible risks. Your healthcare providers can talk with you about using dextroamphetamine and what treatment is best for you. Be sure to talk to your healthcare provider about all of your breastfeeding questions.

If a male takes dextroamphetamine, could it affect fertility (ability to get partner pregnant) or increase the chance of birth defects?

Studies have not been done to see if dextroamphetamine could affect male fertility or increase the chance of birth defects. In general, exposures that fathers or sperm donors have are unlikely to increase risks to a pregnancy. For more information, please see the MotherToBaby fact sheet Paternal Exposures at <https://mothertobaby.org/fact-sheets/paternal-exposures-pregnancy/>.

Please click [here](#) to view references.

Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://mothertobaby.org).

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). *OTIS/MotherToBaby encourages inclusive and person-centered language. While our name still contains a reference to mothers, we are updating our resources with more inclusive terms. Use of the term mother or maternal refers to a person who is pregnant. Use of the term father or paternal refers to a person who contributes sperm.* Copyright by OTIS, January 1, 2022.