

# Lidocaine

---

This sheet is about exposure to lidocaine in pregnancy and while breastfeeding. This information should not take the place of medical care and advice from your healthcare provider.

## ***What is lidocaine?***

Lidocaine is a local anesthetic. Local anesthetics are used to numb areas of the body for short periods of time. Lidocaine has been used as an injection (given by shot), intravenously (by I.V.) and topically (rubbed on the body). Lidocaine can be found in some over-the-counter pain-relieving creams and patches. Lidocaine I.V. has also been used to treat ventricular arrhythmia (abnormal heart rhythm). Injected lidocaine has also been used in some medical or dental procedures or as a therapeutic nerve block to numb a part of the body and relieve pain. Lidocaine is sometimes injected into spinal fluid or the space around the spinal cord ('epidural') to provide pain relief during labor or for a surgical delivery ('cesarian section').

Lidocaine is also found in a topical cream called EMLA®. EMLA® cream also contains the medication prilocaine. For more information on prilocaine, please see the MotherToBaby fact sheet at: <https://mothertobaby.org/fact-sheets/prilocaine>.

## ***I take lidocaine. Can it make it harder for me to get pregnant?***

Studies have not been done to see if lidocaine could make it harder to get pregnant. An experimental animal study did not find that lidocaine would affect fertility (ability to get pregnant).

## ***Does taking lidocaine increase the chance for miscarriage?***

Miscarriage is common and can occur in any pregnancy for many different reasons. Studies have not been done to see if lidocaine increases the chance for miscarriage.

## ***Does taking lidocaine increase the chance of birth defects?***

Every pregnancy starts out with a 3-5% chance of having a birth defect. This is called the background risk. One study involving lidocaine as a local anesthetic did not find a higher chance for birth defects. Also, human case reports and experimental animal studies do not suggest that lidocaine would significantly increase the chance of birth defects.

## ***Does taking lidocaine in pregnancy increase the chance of other pregnancy related problems?***

There are a few case reports of side effects in newborns of pregnant women who received lidocaine at the time of delivery. The infants in these case reports needed treatment for symptoms that included low muscle tone ("floppy"), dilated pupils, difficulty breathing, apnea, and/or seizures. There are also case reports without health concerns in the newborn when lidocaine was used at the time of delivery.

## ***Does taking lidocaine in pregnancy affect future behavior or learning for the child?***

Studies have not been done to see if lidocaine can cause behavior or learning issues for the child.

## ***Breastfeeding while taking lidocaine:***

Lidocaine can get into breastmilk at low levels. However, when swallowed, it is not well absorbed by the baby. Breastfeeding after receiving lidocaine is unlikely to cause problems for a nursing child. Be sure to talk to your healthcare provider about all your breastfeeding questions.

## ***If a man takes lidocaine, could it affect fertility (ability to get a woman pregnant) or increase the chance of birth defects?***

Studies have not been done to see if lidocaine could affect fertility. In general, exposures that men have are unlikely to increase the risks to a pregnancy. For more information, please see the MotherToBaby fact sheet Paternal Exposures at <https://mothertobaby.org/fact-sheets/paternal-exposures-pregnancy/>.

Please click **here** to view references.

**Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://www.MotherToBaby.org).**

---

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, May 1, 2023.