

Listeria Infection (Listeriosis)

This sheet is about having **Listeria** infection (listeriosis) in a pregnancy or while breastfeeding. This information is based on available published literature. It should not take the place of medical care and advice from your healthcare provider.

What is listeriosis?

Listeriosis is an infection caused by the bacteria **Listeria monocytogenes**, often called **Listeria**. Listeriosis is typically caused by eating food that has been contaminated with **this bacteria**, particularly dairy products, leafy vegetables, fish, and meats. **Listeria** can be found everywhere, including your home, restaurants, and other places such as grocery stores and food processing plants. Food with **Listeria** can introduce the infection into the refrigerator where it can spread to other foods. **Listeria** can continue to live in cold temperatures, such as in the refrigerator, but the chance for spreading can be slowed if the refrigerator is kept at a temperature of 40°F / 4°C or lower.

Some foods that are more likely to be contaminated with **Listeria** include unpasteurized (raw) milk, and raw milk products, uncooked meat and fish, uncooked vegetables, deli meat, and soft cheeses. **Listeria** outbreaks can also be found on fruits, like cantaloupes and peaches.

Who is at risk of being affected by listeriosis?

The people most at risk of being affected by listeriosis include women who are pregnant, young children, adults over the age of 60, and people with weakened immune systems. Following some simple food safety guidelines can reduce the chance of getting and spreading listeriosis.

What precautions should I take to lower the chance of getting listeriosis?

To lower the chance of getting and spreading listeriosis and other food-borne illness in **all people**:

- Do not drink unpasteurized milk (also called raw milk) or eat any foods made with unpasteurized milk, including yogurt and ice cream.
- Thoroughly cook raw foods from animal sources.
- Heat foods to steaming (at least 165°F / 74°C) to kill the bacteria.
- Wash raw vegetables and fruit, even if you plan to peel them (remove skin).
- Separate uncooked meats from cooked meats and vegetables, especially with cutting boards to avoid cross-contamination.
- Wash your hands, cutting boards, knives, counters, and sinks well after contact with uncooked foods with a new towel or paper towel.
- Consume ready-to-eat foods as soon as possible.
 - Avoid the "Danger Zone" (temperatures within 40°F and 140°F) when storing foods. Do not leave food out of the refrigerator for more than 2 hours.
- Keep your refrigerator at or below 40°F / 4°C.
- Clean your refrigerator regularly.

Women who are pregnant should take extra precautions to lower the chance of getting listeriosis:

- Do not eat soft cheeses (such as feta, Brie, Camembert, blue-veined cheeses, and Mexican-style cheeses such as queso blanco, queso fresco and Panela) unless they have labels stating that they are made from pasteurized milk. Note: some Mexican-style cheeses made from pasteurized milk have been a source of **Listeria** infections, possibly due to the cheese making process.
- Reheat any leftovers, ready-to-eat foods, hot dogs, cold cuts, deli meat, frozen vegetables, and frozen prepared foods to steaming (at least 165°F / 74°C),

- Prevent getting the juice of deli meats and hot dogs on other foods and surfaces. Wash your hands well after handling deli meats and hot dogs.
- Do not eat refrigerated hummus, pâté, meat spreads or refrigerated smoked seafood, unless it is an ingredient in a fully cooked dish (like a casserole).
- Avoid ready-to-eat salads.

How can I find out if I am infected with Listeria?

Not everyone infected with **Listeria** will develop symptoms. Symptoms could include diarrhea, fever, muscle pain, joint pain, headache, stiff neck, backache, chills, sensitivity to bright light, and/or sore throat with fever and swollen glands. These symptoms can begin days to weeks after eating contaminated food. A blood test can confirm whether you have been infected with listeriosis. If you have eaten contaminated food and do not have symptoms, some experts feel no special testing or treatment is needed. If you think you have listeriosis, talk with your healthcare provider.

I have listeriosis. Can it make it harder for me to become pregnant?

It is not known if listeriosis can make it harder to get pregnant.

Does having / getting listeriosis increase the chance for miscarriage?

Miscarriage is common and can occur in any pregnancy for many different reasons. There is an increased chance of miscarriage when listeriosis occurs in the first trimester of pregnancy. Infections after the first trimester are associated with a lower chance of pregnancy loss.

Does having / getting listeriosis increase the chance of birth defects?

Every pregnancy starts out with a 3-5% chance of having a birth defect. This is called the background risk. Based on the studies reviewed, it is not known if listeriosis increases the chance for birth defects above the background risk.

Would having / getting listeriosis increase the chance of other pregnancy-related problems?

Pregnancies affected by or exposed to **Listeria** can have an increased chance of infection in the uterus, preterm delivery (birth before week 37) or low birth weight (weighing less than 5 pounds, 8 ounces (2500 grams) at birth. Severe infection can increase the chance of stillbirth.

Listeriosis during pregnancy can also increase the chance for serious health problems after birth. Newborn babies infected with **Listeria** can develop either early-onset or late-onset listeriosis. Early onset listeriosis develops 1-2 days after birth, and the baby often has signs of a serious bacterial infection. Late-onset listeriosis occurs 1-2 weeks after birth and usually includes symptoms of meningitis (a condition that causes areas of swelling around the brain and spinal cord in the baby). If not treated quickly, meningitis can lead to long-term problems for some children. Late-onset listeriosis is most likely related to **Listeria** present in the mother's birth canal. Not all babies who are exposed to listeriosis during pregnancy will have problems.

How can I find out if my pregnancy has been affected by Listeria?

An ultrasound can be used to check for an enlarged heart, thickened bowel, and increased thickness of the stomach walls, which may occur in some babies infected with **Listeria**. A blood test can also be performed on the baby after birth to detect whether the baby has been infected with **Listeria**. ***If you suspect you have been infected with Listeria, talk to your healthcare provider as soon as possible.***

Does having / getting listeriosis in pregnancy affect future behavior or learning for the child?

Studies have not been done to see if listeriosis can cause behavior or learning issues for the child.

Breastfeeding while I have listeriosis:

It is not clear if **Listeria** passes through breastmilk. If you have been diagnosed with listeriosis and are breastfeeding, talk to your healthcare provider or your child's pediatrician. Be sure to talk to your healthcare provider about all of your breastfeeding questions.

If a man has listeriosis, can it make it harder to get a woman pregnant or increase the chance of birth defects?

Studies have not been done to see if listeriosis could affect a man's fertility or increase the chance of birth defects. In general, exposures that men have are unlikely to increase risks to a pregnancy. For more information, please see the MotherToBaby fact sheet on Paternal Exposures at <https://mothertobaby.org/fact-sheets/paternal-exposures-pregnancy/>.

Where can I find an updated list of listeria outbreaks?

The Centers for Disease Control and Prevention, better known as the CDC, has a list of all the outbreaks collected since 2011 in the United States. You can access a detailed list of new and previous outbreaks at: <https://www.cdc.gov/listeria/outbreaks/index.html>.

Please click [here](#) for references.

Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://mothertobaby.org).

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, January 1, 2024.