

Lorazepam

This sheet is about exposure to lorazepam in pregnancy and while breastfeeding. This information is based on available published literature. It should not take the place of medical care and advice from your healthcare provider.

What is lorazepam?

Lorazepam is a medication that has been used to treat anxiety and insomnia (trouble falling and/or staying asleep). It has also been used to treat seizures and alcohol withdrawal syndrome. Some brand names for lorazepam are Ativan® and Loreev®. Lorazepam belongs to the class of medication called benzodiazepines.

Sometimes when women find out they are pregnant, they think about changing how they take their medication, or stopping their medication altogether. However, it is important to talk with your healthcare providers before making any changes to how you take your medication. Studies have shown that when anxiety is left untreated during pregnancy, there can be a higher chance for pregnancy complications such as preterm delivery and/or low birth weight. MotherToBaby has a fact sheet on anxiety at <https://mothertobaby.org/fact-sheets/anxiety-fact/>.

Some people have physical symptoms (called withdrawal) when they suddenly stop taking lorazepam. It is not known if or how withdrawal might affect a pregnancy. Your healthcare providers can talk with you about the benefits of treating your condition and the risks of untreated illness during pregnancy.

I take lorazepam. Can it make it harder for me to get pregnant?

Studies have not been done to see if lorazepam can make it harder to get pregnant.

Does taking lorazepam increase the chance of miscarriage?

Miscarriage is common and can occur in any pregnancy for many different reasons. It is not known if lorazepam could increase the chance of miscarriage. Some studies looking at benzodiazepines as a group suggest an increased chance of miscarriage. As there can be many causes of miscarriage, it is hard to know if a medication, the medical condition being treated, or other factors are the cause of a miscarriage.

Does taking lorazepam increase the chance of birth defects?

Birth defects can happen in any pregnancy for different reasons. Out of all babies born each year, about 3 out of 100 (3%) will have a birth defect. It is unlikely that lorazepam significantly increases the chance of birth defects. One study found a possible association with anal atresia (bottom of the intestinal tract is closed off), and another study found an increased chance of pulmonary valve stenosis (abnormal development of the baby's heart). However, other studies did not find an increased chance of birth defects with the use of lorazepam.

Does taking lorazepam in pregnancy increase the chance of other pregnancy-related problems?

Some studies have suggested a higher chance of preterm delivery (birth before week 37) and low birth weight (weighing less than 5 pounds, 8 ounces [2500 grams] at birth) in infants that were exposed to lorazepam in the second half of pregnancy. However, not all studies found a higher chance for these pregnancy-related complications. It is possible that other factors, not lorazepam, caused these complications.

I need to take lorazepam throughout my entire pregnancy. Will it cause withdrawal symptoms in my baby after birth?

The use of lorazepam during pregnancy can cause temporary symptoms in newborns soon after birth. These symptoms are sometimes referred to as withdrawal. Symptoms can include irritability, crying, sleep disturbances, tremors, jitteriness, trouble breathing, or muscle weakness. Not all babies exposed to lorazepam will have these symptoms. If symptoms develop, they usually go away within a few weeks and are not known to have any long-term effects for the baby. It is important that your healthcare providers know you are taking lorazepam so that if symptoms occur your baby can get the care that is best for them.

Does taking lorazepam in pregnancy affect future behavior or learning for the child?

Studies have not been done to see if lorazepam can increase the chance of behavior or learning issues for the child.

Breastfeeding while taking lorazepam:

Lorazepam gets into breast milk in small amounts. No side effects were found in studies of children exposed through breast milk. If you suspect the baby has any symptoms (being too sleepy, poor feeding, or poor weight gain), contact the child's healthcare provider. Be sure to talk to your healthcare provider about all your breastfeeding questions.

If a man takes lorazepam, can it affect fertility or increase the chance of birth defects?

When a male takes lorazepam, it is not expected to affect fertility (ability to get a woman pregnant) or increase the chance of birth defects. In general, exposures that men have are unlikely to increase risks to a pregnancy. For more information, please see the MotherToBaby fact sheet on Paternal Exposures at <https://mothertobaby.org/fact-sheets/paternal-exposures-pregnancy/>.

Please click here for references.

Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://mothertobaby.org).

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, April 1, 2024.