

# Malaria

---

This sheet is about having malaria in a pregnancy or while breastfeeding. This information is based on available published literature. It should not take the place of medical care and advice from your healthcare provider.

## ***What is malaria?***

Malaria is an infectious disease found in parts of the world with warmer weather, especially in tropical and subtropical areas like sub-Saharan Africa, New Guinea, South Asia, Central and South America.

Symptoms of malaria can range from mild to severe. Most people with malaria have fever and flu-like illness with chills, headache, muscle soreness, and extreme tiredness (fatigue). Some people may also have nausea, vomiting, diarrhea, anemia (low red blood cell count), or jaundice (yellowing of the skin and eyes). More rarely, malaria infection may lead to kidney failure, seizures, confusion, coma, or death. Malaria symptoms usually start between 7 to 30 days after infection but can happen up to 1 year after exposure.

## ***How do you get malaria?***

People can develop malaria from being bitten by a mosquito infected with malaria parasites (Plasmodium). Less commonly, malaria infection can come from blood transfusions, organ transplants, or the shared use of needles or syringes contaminated with infected blood. A woman who is pregnant and has malaria may also pass malaria to the baby before or during delivery.

Malaria is not passed through casual contact because it is found only in blood. You cannot get malaria from holding hands or sitting next to someone with malaria. It is not passed like the common cold or flu through coughing or sneezing.

## ***What can I do to prevent getting malaria?***

Since no method of malaria prevention works completely, the Centers for Disease Control and Prevention (CDC) recommend that women who are pregnant should not travel to areas where malaria is found. If it is not possible to avoid travel, ways to help prevent malaria infection include properly using insect repellent (see the MotherToBaby fact sheets on DEET: <https://mothertobaby.org/fact-sheets/deet-nn-ethyl-m-toluamide-pregnancy/> and Insect Repellents: <https://mothertobaby.org/fact-sheets/insect-repellents/>), sleeping in mosquito-free areas, wearing long sleeves and pants, and taking medication to prevent infection before, during, and after travel. The risks associated with malaria infection are likely a greater threat to women who are pregnant and the fetus than any risks from some of the medication used for prevention. If you have questions about a specific medication, talk with your healthcare provider or a MotherToBaby specialist.

## ***I have malaria. Can it make it harder for me to become pregnant?***

It is not known if having malaria can make it harder to get pregnant.

## ***Does having/getting malaria increase the chance for miscarriage?***

Miscarriage is common and can occur in any pregnancy for many different reasons. If a woman has malaria during pregnancy, there is a higher chance of miscarriage.

## ***Does having/getting malaria increase the chance of birth defects?***

Every pregnancy starts out with a 3-5% chance of having a birth defect. This is called the background risk. Malaria is not expected to increase the chance for birth defects above the background risk.

## ***Would having/getting malaria increase the chance of other pregnancy related problems?***

Having malaria during pregnancy can cause a higher chance for preterm birth (birth before 37 weeks of pregnancy), stillbirth, and growth problems in the baby. Symptoms of malaria in the woman who is pregnancy, such as fever, low oxygen levels, or low blood sugar, may also raise the chance of pregnancy complications.

When a woman who is pregnant is infected with malaria, there is a chance that the placenta or baby will become

infected. Infection of the placenta is more common than infection of the baby. Placental infection may prevent the fetus from getting proper amounts of oxygen and nutrients. Infection may also raise the chance for dangerously high blood pressure in the woman who is pregnant. If a baby was infected with malaria during pregnancy, the baby might have symptoms of fever, irritability, feeding problems, breathing problems, sluggishness, paleness, anemia, an enlarged liver and spleen, jaundice, and/or diarrhea in the weeks after birth.

***Does having/getting malaria in pregnancy affect future behavior or learning for the child?***

Some studies have tried to see if malaria can cause behavior or learning issues for a child who was exposed during a pregnancy, but data is very limited. One study showed that malaria infections in pregnancy, especially in late pregnancy, could be related to motor delay (rolling over, crawling, walking). Overall, it is not known if malaria increases the chance for behavior or learning issues.

***What screenings or tests are available to see if the fetus has growth issues?***

Prenatal ultrasounds can be used to monitor the growth of the pregnancy. Talk with your healthcare provider about any prenatal screenings or testing that are available to you. There are no tests available during pregnancy that can tell how much effect there could be on future behavior or learning.

***Breastfeeding while I have malaria:***

Malaria is not passed through breast milk, so breastfeeding will not give your baby malaria. Some medications used to treat malaria might enter the breast milk. Talk with your healthcare provider or contact a MotherToBaby specialist with questions about your specific medication.

One medication to use with caution while breastfeeding is called primaquine. This drug is able to treat malaria infections very well, but it may cause serious red blood cell problems in adults and infants / children who have a genetic condition called glucose-6-phosphate dehydrogenase deficiency (G6PD deficiency). People who need primaquine should be tested for G6PD deficiency before this medication is used. Be sure to talk to your healthcare provider about all of your breastfeeding questions.

***If a man has malaria, can it make it harder to get a partner pregnant or increase the chance of birth defects?***

Studies have not been done to see if malaria could affect a man's fertility (ability to get a woman pregnant) or increase the chance of birth defects. There is one case report of a man who developed low sperm count while ill with malaria. Studies were not done to confirm if malaria was the cause of the reported temporary low sperm count. In general, exposures that men have are unlikely to increase the risks to a pregnancy. For more information, please see the MotherToBaby fact sheet Paternal Exposures at <https://mothertobaby.org/fact-sheets/paternal-exposures-pregnancy/>.

**Please click here for references.**

**Questions? Call 866.626.6847 | Text 855.999.3525 | Email or Chat at [MotherToBaby.org](https://www.MotherToBaby.org).**

---

Disclaimer: MotherToBaby Fact Sheets are meant for general information purposes and should not replace the advice of your health care provider. MotherToBaby is a service of the non-profit Organization of Teratology Information Specialists (OTIS). Copyright by OTIS, November 1, 2023.