



MotherToBaby

Medications & More During Pregnancy & Breastfeeding
Ask The Experts

Fact Sheet

by the **Organization of Teratology Information Specialists (OTIS)**

For more information about us or to find a service in your area,
call **(866) 626-6847**. Visit us online at **www.MotherToBaby.org**.

Find us! Facebook.com/MotherToBaby or @MotherToBaby on Twitter

Meclizine

In every pregnancy, a woman starts out with a 3-5% chance of having a baby with a birth defect. This is called her background risk. This sheet talks about whether exposure to meclizine may increase the risk for birth defects over that background risk. This information should not take the place of medical care and advice from your health care provider.

What is meclizine?

Meclizine is a medication used to treat nausea and vomiting, motion sickness, and vertigo (a feeling of whirling and loss of balance). The medication is an antihistamine (medication used to stop some allergic or cold symptoms) and might cause drowsiness. Some brand names for meclizine have included: Antivert[®], Bonine[®] and Meclicot[®]. Meclizine comes in several over-the-counter (OTC) products such as some Dramamine[®] products.

I just found out I am pregnant. Should I stop taking meclizine?

You should always talk with your healthcare provider before making any changes in your medications. It is important to consider the benefit of your treatment. For more information on nausea and vomiting in pregnancy, please see the MotherToBaby Fact Sheet Nausea and Vomiting in Pregnancy at <https://mothertobaby.org/fact-sheets/nausea-vomiting-pregnancy-nvp/pdf/>.

Can use of meclizine during pregnancy cause birth defects?

Based on the data available, meclizine does not seem to cause an increased chance for birth defects when used in pregnancy.

Can use of meclizine cause other pregnancy complications?

There are no data looking at the use of meclizine and the chance for other complications to a pregnancy.

Can I take meclizine while breastfeeding?

There is limited information on the use of meclizine while breastfeeding. Occasional doses of meclizine while breastfeeding are unlikely to cause a problem for your baby. However, if you need to use this medicine often or in large doses, talk with your healthcare provider. It is possible, but not proven, that antihistamines in general may lower the amount of milk a woman makes. This might be more likely to occur if antihistamines are used in combination with an oral decongestant like pseudoephedrine (Sudafed[®]), or if used before breastfeeding is fully established (when baby is first born). Be sure to talk to your healthcare provider about all of your breastfeeding questions.

What if the father of the baby takes meclizine?

There are no studies looking at how a man's use of meclizine would affect his fertility or a partner's pregnancy. In general, exposures that fathers have are unlikely to increase risks to a pregnancy. For more information, please see the MotherToBaby fact sheet Paternal Exposures and Pregnancy at <https://mothertobaby.org/fact-sheets/paternal-exposures-pregnancy/pdf/>.

References:

- Gilboa SM, et al. 2009. National Birth Defects Prevention Study: Use of antihistamine medications during early pregnancy and isolated major malformations. Birth Defects Res A Clin Mol Teratol 85(2):137-150.
- Ito S, et al. 1993. Prospective follow-up of adverse reactions in breast-fed infants exposed to maternal medication. Am J Obstet Gynecol; 168:1393-9.

- Kallen B, Mottet I. 2003. Delivery outcome after the use of meclizine in early pregnancy. *Eur J Epidemiol* 18:665-669.
- King CT, et al. 1965. Antihistamines and teratogenicity in the rat. *J Pharmacol Exp Ther*; 147:391-8.
- Lenz W. 1966. Malformations caused by drugs in pregnancy. *Am J Dis Child*; 112:99-106.
- Messinis IE, et al. 1985. Histamine H1 receptor participation in the control of prolactin secretion in postpartum. *J Endocrinol Investig*;8:143-6
- Michaelis J, et al. 1983. Prospective study of suspected associations between certain drugs administered during early pregnancy and congenital malformations. *Teratology* 27:57-64.
- Wilk AL, et al. 1978. Chlorcyclizine induction of cleft palate in the rat: degradation of palatal glycosaminoglycans. *Teratology*; 18(2):199-209.

January, 2018