

Fetal Development Poster

16" x 24"

Critical periods of your baby's development



WHAT ARE STRUCTURAL BIRTH DEFECTS?

Structural birth defects are abnormalities in the structure of body parts. Examples include cleft palate, heart defects and spina bifida.



WHAT ARE FUNCTIONAL BIRTH DEFECTS?


Functional birth defects are problems in how a body system works. Examples include hearing and vision loss.



WHAT CAUSES BIRTH DEFECTS?

Birth defects can be caused by genetic (mutations) or environmental factors (teratogens). Examples of teratogens include alcohol, certain medications, and infections.

This chart shows the most sensitive times of a baby's development during the 38 weeks of pregnancy*



*This fetal chart shows the 38 weeks of pregnancy. Since it is difficult to know exactly when conception occurs, health care providers calculate a woman's due date 40 weeks from the start of her last menstrual cycle. Adapted from Moore, 1993 and the National Organization on Fetal Alcohol Syndrome (NOFAS) 2009.



Hablamos Español

ASK THE EXPERTS

about the safety of medications and other exposures during pregnancy.

Call 877.311.8972

Visit mothertobaby.org

Download App



MotherToBaby
PREGNANCY STUDIES

RESEARCH FUELED BY MOMS

For a list of our current observational studies and to see if you qualify, visit mothertobaby.org/studies